

**Bold type** indicates authors of commentaries, editorials, and Resource Corners; (No.) indicates issue number.

- Abramson MJ, 43  
 Akl E, A-15 (No. 1)  
 Alkhenizan A, 64  
 Alonso JJ, 38  
 Anavekar NS, 51  
 Andersen J, 10  
 Anderson M, A-13 (No. 2)  
 Andersson SO, 24  
 Andr  n O, 24  
 Annane D, 30  
 Armstrong D, 6  
 Avenell A, 18
- Bach RG, 3  
 Banks SM, 31  
 Barry MJ, 24  
 Bates ER, 3  
 Baughman KL, 57  
 Beecher LH, 4  
**Begh   C, 8**  
 Beglinger C, 78  
 Bellissant E, 30  
 Bell-Syer SE, 8  
 Belshe RB, 68  
 Ben-Dor I, 32  
 Benowitz NL, 12  
**Bentz CL, 12**  
 Berghold A, 63  
 Betteridge DJ, 29  
 Bingham S, 52  
 Bjelakovic G, 20  
 Blum AL, 78  
 Boetes C, 23  
 Bollaert PE, 30  
 Bonino F, 34  
 Booker L, 14  
**Borzak S, 39**  
 Braunwald E, 66  
 Brekelmans CT, 23  
 Bridges JF, 35  
 Bronson DL, 16  
**Brott B, 38**  
 Brown MD, 77  
 Brown TJ, 75  
 Burmester GR, 46  
 Byrne CD, 18-19
- Campbell IA, 44  
 Cannon CP, 3  
 Cannon J, 68  
 Capece R, 16  
 Cardozo L, 48  
 Carlberg B, 59  
 Castro-Beiras A, 38  
 Catalano C, 49  
 Cepeda-Benito A, 13  
 Chan FKL, 75, 76  
 Chaves PH, 81  
 Chertow GM, 50  
 Chessman AW, 21  
 Chiba N, 6
- Chirchilli VM, 43  
 Choudhury A, 58  
 Chronicle E, 21  
 Chu K, 22  
**Chung RT, A-12 (No. 1), 10**  
 Coccheri S, 61  
 Coffey CS, 36  
 Cohen AT, 40  
**Cohrsen A, A-13 (No. 2)**  
 Colhoun HM, 29  
 Colman I, 77  
 Conaway MR, 17  
 Cook D, 5  
 Corsello A, 35  
 Crowther JQ, 17  
 Cruickshank JK, 59  
 Cullen L, 22  
 Cullum N, 8
- de Jong Z, 73  
 Deans KJ, 31  
 Deeks JJ, 70  
 Delaney BC, 74  
 Demicheli V, 70  
**Denman AM, 76**  
 Desai AS, 57  
 Domanski MJ, 66  
 Donowitz GR, 33  
 Dooley DP, 45  
 Dorian P, 58  
 Douglas JG, 44  
 Dunlap ME, 60  
**Dunn A, 71**  
 Durrington PN, 29
- Edmonds M, 42**  
 Eichacker PQ, 31  
 Elliott R, 75  
 Erath S, 13  
**Eva KW, A-8-A-9 (No. 3)**  
 Ezekowitz JA, 35
- Fan D, 50  
 Fang JC, 57  
 Farkouh ME, 46  
 Fassi A, 65  
**Fernandes CMB, 4**  
**Fernandez-Avil  s F, 38**  
 Fischer TL, 4  
**Fletcher RH, 20**  
 Ford JG, 43  
 Forman D, 74  
 Fowler SE, 66  
 Francis RM, 44  
 Frech SA, 69  
 Freedman M, 15  
 Freudenberg RS, 64  
 Fried M, 78  
**Furberg CD, 1**  
 Furlong NJ, 62
- Gami AS, 29  
 Garcia DA, 7  
 Garg AX, 50-51  
 Ghosh AK, A-12 (No. 2)  
 Gill SS, 15  
**Glasziou P, A-8-A-10 (No. 2)**  
**Glatt AE, 33**  
 Glenn GM, 69  
 Gluud C, 20  
 Go AS, 50  
 Goldberg AC, 16  
 Goudswaard AN, 62  
 Gourlay SG, 12  
**Granger CB, 66**  
 Greyber E, 1  
 Guralnik JM, 81
- Hagens VE, 53  
 Hallstrom AP, 2  
**Hankey G, 41**  
 Hannonen P, 9  
 Harrington RA, 46  
 Harrison RA, 72  
 Hart L, 47  
**Haynes RB, A-8-A-9 (No. 1), A-8-A-10 (No. 2)**  
 Hebert PR, 36  
 Hepburn MJ, 45  
**Hernandez AF, 57**  
**Hicks NR, A-10 (No. 1)**  
**Hillegeass WB, 38**  
**Hirsch C, 14-15**  
 Hohnloser SH, 58  
 Holdgate A, 42  
 Hooper L, 72, 75  
 Hsu C, 50  
 Hughes JR, 67  
 Hung LC, 76  
**Hunt D, 65**  
 Huszti E, 35  
**Hyde C, A-11 (No. 2)**
- Iannaccone R, 49  
 Ilieva AP, 65  
 Innes GD, 77  
 Israel E, 43  
**Jamal SA, 44**
- Jefferson TO, 70  
 Johansson JE, 24  
 Juurlink DN, 25
- Kaul P, 35  
**Kendall DN, 62-63**  
 Kenney RT, 69  
 Kessels AG, 80  
 Khaw KT, 52  
 Kirshner H, 46  
**Klink K, A-13 (No. 2)**  
 Ko DT, 36  
 Kopp A, 25

- Koretz RL, A-13 (No. 1), 10-11  
Korpela M, 9  
Kostis JB, 64  
Kriege M, 23  
Krishnan KR, 79  
Kuck KH, 58  
Kudva Y, 65  
Kuntz RE, 41
- Laasonen L, 9  
Laghi A, 49  
Lancaster T, 13, 67  
Langley JM, 68-69  
Lau DC, 19  
Lau GK, 34  
Lauritsen K, 6  
Laville M, 25  
Lee PE, 15, 79  
Legood R, 8  
Leiter LA, 17  
Leizorovitz A, 40  
Lensing AW, 7, 71  
Lesbros-Pantoflickova D, 78  
Li SK, 19  
Lindholm LH, 59  
Lip GY, 58  
Liu J, 16  
Loder EW, 77  
Lose G, 48  
Luckmann R, 23
- Maisel WH, 57  
Majumdar SR, 60  
Mamdani MM, 25  
Marcellin P, 34  
Marubini E, 61  
Massie BM, 37  
Maxwell L, 73  
Mayo-Smith MF, 4  
McAlister FA, 35  
McClish D, 48  
McCulloch CE, 50  
McInnes E, 8  
McMurray JJ, 51  
Meade M, 30-31  
Michetti P, 78  
Minneci PC, 31  
Mitchel YB, 16  
Moayyedi P, 74  
Morris AM, 45  
Mueller C, A-14-A-15 (No. 1)  
Muenz LR, 69  
Mulleners W, 21  
Munneke M, 73  
Murali NS, A-12 (No. 2)  
Mysler E, 46
- Natanson C, 31  
Nelemans PJ, 80  
Neri Serneri GG, 61  
Newman D, 53
- Newman FK, 68  
Nichol G, 35  
Nidiry JJ, 34  
Nikolova D, 20  
Norman GR, A-8-A-9 (No. 3)  
Norris SL, 18
- O'Malley OG, 36  
O'Flaherty M, 81  
Ormiston T, 1  
Ornato JP, 2
- Padwal R, 19  
Paul M, 32  
Pfeffer MA, 60  
Pfisterer M, 39  
Plank J, 63  
Pollock T, 42  
Prandoni P, 7  
Prescott RJ, 44  
Primack BA, 78  
Prins MH, 7, 71  
Prochazka AV, 67  
Psaty BM, 1
- Qushmag I, 5
- Raina P, 14  
Ransohoff DE, 49  
Reid DM, 44  
Reynoso JT, 13  
Rivetti D, 70  
Roberts I, 5  
Rochon P, 79  
Rockstroh JK, 11  
Rodriguez-Torres M, 11  
Roose SR, 79  
Ruggenenti P, 65  
Rutten GE, 62
- Sackeim HA, 79  
Sacks HS, 70  
Salpeter EE, 1  
Salpeter SR, 1  
Samuelsson O, 59  
Sander O, 9  
Sandercock P, 5  
Santaguida P, 14  
Sapre A, 16  
Schünemann HJ, A-15 (No. 1)  
Schatz IJ, A-14 (No. 2)  
Schnitzer TJ, 46  
Schwam E, A-14 (No. 1)  
Siebenhofer A, 63  
Sigal RJ, 52  
Simon JA, 72  
Simonetti RG, 20  
Skidmore PJ, 45  
Smith DA, 61  
Smith SA, 29  
Spandorfer J, 40
- Stead LF, 12, 67  
Stolk RP, 62  
Suen BY, 76  
Sullivan F, A-11 (No. 1)
- Talley NJ, 6, 74  
Taylor AL, 37  
Taylor D, 22  
Taylor S, 22  
TenVergert EM, 53  
Teppler H, 33  
Thacker HL, 48  
Thompson RL, 72  
Torriani FJ, A-12 (No. 1), 11  
Tugwell P, 73  
Turpie AG, 40
- Vakil N, 74  
Valk GD, 62  
van den Belt AG, 71  
van Dongen CJ, 71  
Vasbinder GB, 80  
Velazquez EJ, 51  
Veldhuyzen Van Zanten SJ, 74  
Vermeulen KM, 53  
Versi E, 48  
Vidal L, 32  
Villar CP, 69  
Violi F, 61  
Volberding P, 10
- Walker-Dilks C, A-10 (No. 3)  
Walsh JM, 1  
Walsh TJ, 33  
Wareham N, 52  
Weintraub WS, 3  
Weise WJ, 80  
Weisfeldt M, 2  
Welborn J, 32  
Werre SR, A-10 (No. 3)  
Wholey MH, 41  
Wiebe N, 35  
Wilczynski N, A-8-A-9 (No. 1)  
Wild S, 18-19  
Wilson AC, 64  
Wolf AM, 17  
Wong VWS, 75  
Worster A, 2  
Wyer P, 22
- Xue QL, 81
- Yadav JS, 41  
Yancy C, 37  
Yates D, 5  
Young JB, 60
- Zhang X, 18  
Ziesche S, 37  
Zwiderman AH, 73

**A**

- Administration, inhalation
  - Nebulized lidocaine before nasogastric tube insertion reduced patient discomfort but increased risk for nasal bleeding, 22
- Adrenergic  $\beta$ -agonists
  - Gly/Gly and Arg/Arg genotype responses to albuterol differed in mild asthma, 43
- Adrenergic  $\beta$ -antagonists
  - Review:  $\beta$ -blockers reduced mortality and therapy withdrawal in heart failure but increased dizziness and bradycardia, 36
- African continental ancestry group
  - Isosorbide dinitrate plus hydralazine was effective for advanced heart failure in black patients, 37
- Albuminuria
  - Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65
- Albuterol
  - Gly/Gly and Arg/Arg genotype responses to albuterol differed in mild asthma, 43
- Alcohol withdrawal delirium
  - Review: Sedative-hypnotic agents reduce mortality and duration of delirium in the alcohol withdrawal syndrome, 4
- Amphotericin B
  - Caspofungin was noninferior to amphotericin B for invasive fungal infections in persistent fever and neutropenia and was better tolerated, 33
- Angina
  - An early invasive strategy improved long-term freedom from major adverse cardiac events in older patients with angina, 39
  - An early invasive strategy reduced death or MI better than a conservative strategy in unstable angina and non-STEMI at advanced age, 3
- Angiography
  - Computed tomographic angiography and magnetic resonance angiography were specific, but could not rule out renal artery stenosis, 80
- Angioplasty, transluminal, percutaneous coronary
  - Invasive strategy within 24 hours of thrombolysis reduced death, nonfatal reinfarction, and ischemia-induced revascularization in STEMI, 38
- Angiotensin-converting enzyme inhibitors
  - Candesartan reduced cardiovascular mortality and morbidity in chronic heart failure with low ejection fraction, 60
  - Loop diuretics and angiotensin-converting enzyme inhibitors increased risk for hospitalization for lithium toxicity, 25
  - Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65
  - Trandolapril did not reduce cardiovascular death or other events in stable coronary artery disease, 66
- Antiarrhythmia agents
  - Rate control was more cost-effective than rhythm control in persistent atrial fibrillation, 53
- Antibacterial agents
  - Review: Oral and intravenous antibiotics do not differ for effectiveness in febrile neutropenic patients with cancer, 32
- Antibodies, viral
  - Intradermal injection of reduced-dose influenza vaccine was highly immunogenic in persons  $\leq 60$  years of age but less so in persons  $> 60$  years, 68
  - Intradermal injection of reduced-dose influenza vaccine was immunogenic in young adults, 69
- Anticholesteremic agents
  - Atorvastatin reduced major cardiovascular disease events in type 2 diabetes mellitus, 29
  - Ezetimibe plus simvastatin lowered lipid levels more than simvastatin monotherapy in primary hypercholesterolemia, 16
- Anticoagulants
  - Rate control was more cost-effective than rhythm control in persistent atrial fibrillation, 53
- Anticonvulsants
  - Review: Anticonvulsants are better than placebo for reducing the frequency of migraine attacks, 21
- Antidepressive agents
  - Review: Bupropion and nortriptyline each increase smoking cessation rates, 67
  - Review: Some therapies for treating the irritable bowel syndrome may be effective for symptom improvement, 78
- Antiemetics
  - Review: Intravenous metoclopramide is better than placebo for reducing pain in acute migraine in the emergency department, 77
- Antifungal agents
  - Caspofungin was noninferior to amphotericin B for invasive fungal infections in persistent fever and neutropenia and was better tolerated, 33
- Antihypertensive agents
  - Review: Atenolol may be ineffective for reducing cardiovascular morbidity or all-cause mortality in primary hypertension, 59
- Antiinflammatory agents
  - Review: Misoprostol or COX-2-specific or selective NSAIDs reduce gastrointestinal complications and symptomatic ulcers, 75
  - Review: Nonsteroidal antiinflammatory drugs provide better pain relief than do opioids for acute renal colic, 42
- Antiobesity agents
  - Review: Fluoxetine, orlistat, and sibutramine modestly reduce weight in type 2 diabetes, 18
  - Review: Orlistat and sibutramine are modestly effective for weight loss at 1 year, 19
- Antioxidants
  - Review: Antioxidant supplementation does not reduce gastrointestinal cancer, 20
- Antipsychotic agents
  - Review: Limited evidence supports the use of atypical antipsychotic drugs in behavioral and psychological symptoms of dementia, 15
- Antirheumatic agents
  - Long-term use of combination DMARDs did not sustain disease remissions but delayed joint damage in early rheumatoid arthritis, 9
- Antiviral agents
  - Peginterferon  $\alpha$ -2a alone or combined with lamivudine increased response rates more than lamivudine alone, 34
- Appetite depressants
  - Review: Fluoxetine, orlistat, and sibutramine modestly reduce weight in type 2 diabetes, 18
  - Review: Orlistat and sibutramine are modestly effective for weight loss at 1 year, 19
- Arrhythmia
  - A prophylactically implanted cardioverter defibrillator did not reduce all-cause mortality after a recent myocardial infarction, 58
- Arthritis
  - Celecoxib was noninferior to diclofenac plus omeprazole for preventing gastroduodenal ulcer recurrence in high-risk patients with arthritis, 76
  - High-intensity exercise for rheumatoid arthritis was associated with less joint damage of the hands and feet than physical therapy, 73
  - Long-term use of combination DMARDs did not sustain disease remissions but delayed joint damage in early rheumatoid arthritis, 9

Arthrography

Long-term use of combination DMARDs did not sustain disease remissions but delayed joint damage in early rheumatoid arthritis, 9

Asthma

Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

Gly/Gly and Arg/Arg genotype responses to albuterol differed in mild asthma, 43

Atenolol

Review: Atenolol may be ineffective for reducing cardiovascular morbidity or all-cause mortality in primary hypertension, 59

Atrial fibrillation

Rate control was more cost-effective than rhythm control in persistent atrial fibrillation, 53

Azetidines

Ezetimibe plus simvastatin lowered lipid levels more than simvastatin monotherapy in primary hypercholesterolemia, 16

B

Bandages

Below-knee elastic compression stockings reduced development of the postthrombotic syndrome in proximal deep venous thrombosis, 7

Beds

Review: Foam-based, constant low-pressure mattresses are better than standard hospital mattresses for reducing pressure ulcers, 8

Behavioral symptoms

Review: Limited evidence supports the use of atypical antipsychotic drugs in behavioral and psychological symptoms of dementia, 15

Bone density

Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

Breast neoplasms

Magnetic resonance imaging was more sensitive than mammography for detecting breast cancer in high-risk women, 23

C

Calcium

Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

Calcium-channel blockers

Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65

Candesartan

Candesartan reduced cardiovascular mortality and morbidity in chronic heart failure with low ejection fraction, 60

Candidiasis, vulvovaginal

Review: Vaginal signs and symptoms perform poorly in diagnosing vaginal candidiasis (Letter), A-13 (No. 2)

Review: Vaginal signs and symptoms perform poorly in diagnosing vaginal candidiasis (Correction), A-13 (No. 2)

Cardiac pacing, artificial

Review: Cardiac resynchronization therapy reduces mortality and hospitalization for heart failure, 35

Cardiomyopathies

Review: Implantable cardioverter defibrillators reduce all-cause mortality in nonischemic cardiomyopathy, 57

Cardiopulmonary resuscitation

Volunteers trained in CPR and the use of automated external defibrillators increased survival after out-of-hospital cardiac arrest, 2

Cardiovascular agents

An early invasive strategy improved long-term freedom from major adverse cardiac events in older patients with angina, 39

Cardiovascular disease

Atorvastatin reduced major cardiovascular disease events in type 2 diabetes mellitus, 29

Hemoglobin A<sub>1c</sub> levels were associated with increased cardiovascular disease and all-cause mortality in persons with and without diabetes, 52

Reduced glomerular filtration rate was associated with increased death, cardiovascular events, and hospitalization, 50

Renal impairment increased mortality and cardiovascular complications after myocardial infarction, 51

Review: n-3 fatty acids do not reduce mortality, cardiovascular events, or cancer in patients at risk for cardiovascular disease, 72

Carotid stenosis

Stenting was noninferior to endarterectomy in patients with severe carotid artery stenosis and coexisting conditions, 41

Cellulitis

A 10-day regimen of levofloxacin was not needed in patients with uncomplicated cellulitis who had an acceptable 5-day response, 45

Chlorthalidone

Diuretic-based therapy reduced cardiovascular mortality in older patients with isolated systolic hypertension and diabetes, 64

Citalopram

Citalopram did not differ from placebo for treatment of depression in patients > 75 years of age, 79

Clinical competence

Does clinical experience make up for failure to keep up to date? (Editorial), A-8-A-9 (No. 3)

Clonidine

Review: Clonidine is more effective than placebo for long-term smoking cessation but has side effects, 12

Colic

Review: Nonsteroidal antiinflammatory drugs provide better pain relief than do opioids for acute renal colic, 42

Colonic polyps

Computed tomographic colonography without cathartic preparation performed well in detecting colorectal polyps, 49

Colonography, computed tomographic

Computed tomographic colonography without cathartic preparation performed well in detecting colorectal polyps, 49

Coronary disease

Trandolapril did not reduce cardiovascular death or other events in stable coronary artery disease, 66

Cyclooxygenase inhibitors

Review: Misoprostol or COX-2-specific or selective NSAIDs reduce gastrointestinal complications and symptomatic ulcers, 75

D

Decision making

*Decision Making in Health and Medicine. Integrating Evidence and Values* (Resource Corner), A-11 (No. 2)

From research evidence to context: The challenge of individualizing care (Letter), A-14 (No. 2)

Decubitus ulcer

Review: Foam-based, constant low-pressure mattresses are better than standard hospital mattresses for reducing pressure ulcers, 8

Defibrillators, implantable

Review: Implantable cardioverter defibrillators reduce all-cause mortality in nonischemic cardiomyopathy, 57

A prophylactically implanted cardioverter defibrillator did not reduce all-cause mortality after a recent myocardial infarction, 58

## Dementia

- Review: Limited evidence supports the use of atypical antipsychotic drugs in behavioral and psychological symptoms of dementia, 15
- Review: Pharmacotherapy improves cognitive symptoms in dementia, 14

## Depressive disorder

- Citalopram did not differ from placebo for treatment of depression in patients > 75 years of age, 79

## Diabetes mellitus

- Atorvastatin reduced major cardiovascular disease events in type 2 diabetes mellitus, 29
- A dietitian-led intervention reduced weight and waist circumference in obese patients with type 2 diabetes, 17
- Diuretic-based therapy reduced cardiovascular mortality in older patients with isolated systolic hypertension and diabetes, 64
- Picotamide reduced all-cause mortality more than aspirin in type 2 diabetes mellitus and peripheral arterial disease, 61
- Review: Fluoxetine, orlistat, and sibutramine modestly reduce weight in type 2 diabetes, 18
- Review: Insulin monotherapy and insulin combined with oral hypoglycemic agents provide similar glycemic control, 62
- Review: Short-acting insulin analogues reduce glycosylated hemoglobin more than regular human insulin but only in adults with type 1 diabetes, 63
- Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65

## Diagnostic services

- Screening for Diseases: Prevention in Primary Care* (Resource Corner), A-11 (No. 1)

## Diclofenac

- Celecoxib was noninferior to diclofenac plus omeprazole for preventing gastroduodenal ulcer recurrence in high-risk patients with arthritis, 76

## Dietary fiber

- Review: Some therapies for treating the irritable bowel syndrome may be effective for symptom improvement, 78

## Disabled persons

- A hemoglobin level 140 g/L was associated with a lower mortality risk than levels  $\leq 120$  g/L in older women with disabilities, 81

## Diuretics

- Diuretic-based therapy reduced cardiovascular mortality in older patients with isolated systolic hypertension and diabetes, 64
- Loop diuretics and angiotensin-converting enzyme inhibitors increased risk for hospitalization for lithium toxicity, 25

## Dyspepsia

- Review: Proton-pump inhibitor therapy reduces symptoms in nonulcer dyspepsia better than placebo, 74

## Dyspnea

- Rapid measurement of B-type natriuretic peptides reduced time to discharge and treatment costs in patients with acute dyspnea (Letter), A-14-A-15 (No. 1)

## E

### Education, medical, continuing

- Does clinical experience make up for failure to keep up to date? (Editorial), A-8-A-9 (No. 3)

### Electric countershock

- Rate control was more cost-effective than rhythm control in persistent atrial fibrillation, 53
- Review: Cardiac resynchronization therapy reduces mortality and hospitalization for heart failure, 35
- Volunteers trained in CPR and the use of automated external defibrillators increased survival after out-of-hospital cardiac arrest, 2

## Emergency medical services

- Volunteers trained in CPR and the use of automated external defibrillators increased survival after out-of-hospital cardiac arrest, 2

## Endarterectomy, carotid

- Stenting was noninferior to endarterectomy in patients with severe carotid artery stenosis and coexisting conditions, 41

## Estrogen replacement therapy

- Review: Estrogen improves symptoms of overactive bladder in postmenopausal women, 48
- Review: Hormone replacement therapy may reduce the risk for death in younger but not older postmenopausal women, 1

## Etidronic acid

- Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

## Evidence-based medicine

- Decision Making in Health and Medicine. Integrating Evidence and Values* (Resource Corner), A-11 (No. 2)
- Essential Evidence-based Medicine* (Resource Corner), A-12 (No. 2)
- Evidence-based Public Health* (Resource Corner), A-10 (No. 1)
- Finding the gold in MEDLINE: Clinical Queries (Editorial), A-8-A-9 (No. 1)
- From research evidence to context: The challenge of individualizing care (Letter), A-14 (No. 2)
- The paths from research to improved health outcomes (Editorial), A-8-A-10 (No. 2)
- Weighted event rates (Editorial), A-10 (No. 3)

## Exercise therapy

- High-intensity exercise for rheumatoid arthritis was associated with less joint damage of the hands and feet than physical therapy, 73

## F

### Fatty acids, omega-3

- Review: n-3 fatty acids do not reduce mortality, cardiovascular events, or cancer in patients at risk for cardiovascular disease, 72

### Fever

- Review: Oral and intravenous antibiotics do not differ for effectiveness in febrile neutropenic patients with cancer, 32

### Fluoxetine

- Review: Fluoxetine, orlistat, and sibutramine modestly reduce weight in type 2 diabetes, 18

### Fractures

- Review: Hormone replacement therapy may reduce the risk for death in younger but not older postmenopausal women, 1

### Fractures, spontaneous

- Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

## G

### Gastroesophageal reflux

- Esomeprazole was not better than omeprazole for resolving heartburn in endoscopy-negative reflux disease, 6

### Gastrointestinal neoplasms

- Review: Antioxidant supplementation does not reduce gastrointestinal cancer, 20

### Glomerular filtration rate

- Reduced glomerular filtration rate was associated with increased death, cardiovascular events, and hospitalization, 50
- Renal impairment increased mortality and cardiovascular complications after myocardial infarction, 51



Glucocorticoids

- Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44
- Methylprednisolone caused an increase in death after head injury, 5
- Review: Glucocorticoids reduced mortality in sepsis in recent (post-1997) but not previous (pre-1989) trials, or all trials combined, 31

H

Heart arrest

- Volunteers trained in CPR and the use of automated external defibrillators increased survival after out-of-hospital cardiac arrest, 2

Heart failure, congestive

- Candesartan reduced cardiovascular mortality and morbidity in chronic heart failure with low ejection fraction, 60
- Isosorbide dinitrate plus hydralazine was effective for advanced heart failure in black patients, 37
- Rapid measurement of B-type natriuretic peptides reduced time to discharge and treatment costs in patients with acute dyspnea (Letter), A-14-A-15 (No. 1)
- Review:  $\beta$ -blockers reduced mortality and therapy withdrawal in heart failure but increased dizziness and bradycardia, 36
- Review: Cardiac resynchronization therapy reduces mortality and hospitalization for heart failure, 35

Heartburn

- Esomeprazole was not better than omeprazole for resolving heartburn in endoscopy-negative reflux disease, 6

Hemoglobin

- Hemoglobin A<sub>1c</sub> levels were associated with increased cardiovascular disease and all-cause mortality in persons with and without diabetes, 52
- A hemoglobin level 140 g/L was associated with a lower mortality risk than levels  $\leq 120$  g/L in older women with disabilities, 81

Heparin

- Dalteparin reduced venous thromboembolic events without increased bleeding in acutely ill medical patients, 40
- Review: Low-molecular-weight heparin reduces recurrent venous thromboembolism better than unfractionated heparin, 71

Hepatitis B

- Peginterferon  $\alpha$ -2a alone or combined with lamivudine increased response rates more than lamivudine alone, 34

Hepatitis C

- Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections (Letter), A-12-A-13 (No. 1)

Hydralazine

- Isosorbide dinitrate plus hydralazine was effective for advanced heart failure in black patients, 37

Hypercholesterolemia

- Ezetimibe plus simvastatin lowered lipid levels more than simvastatin monotherapy in primary hypercholesterolemia, 16

Hypertension

- Diuretic-based therapy reduced cardiovascular mortality in older patients with isolated systolic hypertension and diabetes, 64
- Review: Atenolol may be ineffective for reducing cardiovascular morbidity or all-cause mortality in primary hypertension, 59
- Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65

Hypoglycemic agents

- Review: Insulin monotherapy and insulin combined with oral hypoglycemic agents provide similar glycemic control, 62
- Review: Short-acting insulin analogues reduce glycosylated hemoglobin more than regular human insulin but only in adults with type 1 diabetes, 63

I

Ibuprofen

- Lumiracoxib reduced ulcer complications compared with ibuprofen and naproxen in osteoarthritis and did not increase cardiovascular outcomes, 46-47

Indoles

- Trandolapril did not reduce cardiovascular death or other events in stable coronary artery disease, 66

Influenza vaccine

- Review: Vaccination reduces the incidence of serologically confirmed influenza in healthy adults, 70

Influenza

- Intradermal injection of reduced-dose influenza vaccine was highly immunogenic in persons  $\leq 60$  years of age but less so in persons  $> 60$  years, 68
- Intradermal injection of reduced-dose influenza vaccine was immunogenic in young adults, 69
- Review: Vaccination reduces the incidence of serologically confirmed influenza in healthy adults, 70

Inotropic agents

- Review: Pharmacotherapy improves cognitive symptoms in dementia, 14

Insulin

- Review: Insulin monotherapy and insulin combined with oral hypoglycemic agents provide similar glycemic control, 62
- Review: Short-acting insulin analogues reduce glycosylated hemoglobin more than regular human insulin but only in adults with type 1 diabetes, 63

Interferon  $\alpha$ -2a

- Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections, 10
- Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections (Letter), A-12-A-13 (No. 1)
- Peginterferon  $\alpha$ -2a alone or combined with lamivudine increased response rates more than lamivudine alone, 34

Interferon  $\alpha$ -2b

- Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections, 10
- Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections (Letter), A-12-A-13 (No. 1)

Intubation, gastrointestinal

- Nebulized lidocaine before nasogastric tube insertion reduced patient discomfort but increased risk for nasal bleeding, 22

Irritable bowel syndrome

- Review: Some therapies for treating the irritable bowel syndrome may be effective for symptom improvement, 78

Isosorbide dinitrate

- Isosorbide dinitrate plus hydralazine was effective for advanced heart failure in black patients, 37

J

Joints

- High-intensity exercise for rheumatoid arthritis was associated with less joint damage of the hands and feet than physical therapy, 73
- Long-term use of combination DMARDs did not sustain disease remissions but delayed joint damage in early rheumatoid arthritis, 9

K

Kidney diseases

- Reduced glomerular filtration rate was associated with increased death, cardiovascular events, and hospitalization, 50
- Renal impairment increased mortality and cardiovascular complications after myocardial infarction, 51

Review: Nonsteroidal antiinflammatory drugs provide better pain relief than do opioids for acute renal colic, 42

## L

### Lactones

Review: Orlistat and sibutramine are modestly effective for weight loss at 1 year, 19

### Lamivudine

Peginterferon  $\alpha$ -2a alone or combined with lamivudine increased response rates more than lamivudine alone, 34

### Left ventricular dysfunction

Candesartan reduced cardiovascular mortality and morbidity in chronic heart failure with low ejection fraction, 60

### Lidocaine

Nebulized lidocaine before nasogastric tube insertion reduced patient discomfort but increased risk for nasal bleeding, 22

### Lithium

Loop diuretics and angiotensin-converting enzyme inhibitors increased risk for hospitalization for lithium toxicity, 25

## M

### Magnetic resonance angiography

Computed tomographic angiography and magnetic resonance angiography were specific, but could not rule out renal artery stenosis, 80

Magnetic resonance imaging was more sensitive than mammography for detecting breast cancer in high-risk women, 23

### Mammography

Magnetic resonance imaging was more sensitive than mammography for detecting breast cancer in high-risk women, 23

### Mass screening

*Screening for Diseases: Prevention in Primary Care* (Resource Corner), A-11 (No. 1)

### MEDLINE

Finding the gold in MEDLINE: Clinical Queries (Editorial), A-8-A-9 (No. 1)

### Meta-analysis

Weighted event rates (Editorial), A-10 (No. 3)

### Methylprednisolone

Methylprednisolone caused an increase in death after head injury, 5

### Metoclopramide

Review: Intravenous metoclopramide is better than placebo for reducing pain in acute migraine in the emergency department, 77

### Migraine

Review: Intravenous metoclopramide is better than placebo for reducing pain in acute migraine in the emergency department, 77

### Mortality

A hemoglobin level 140 g/L was associated with a lower mortality risk than levels  $\leq 120$  g/L in older women with disabilities, 81

### Mycoses

Caspofungin was noninferior to amphotericin B for invasive fungal infections in persistent fever and neutropenia and was better tolerated, 33

### Myocardial infarction

An early invasive strategy reduced death or MI better than a conservative strategy in unstable angina and non-STEMI at advanced age, 3

Invasive strategy within 24 hours of thrombolysis reduced death, nonfatal reinfarction, and ischemia-induced revascularization in STEMI, 38

Lumiracoxib reduced ulcer complications compared with ibuprofen and naproxen in osteoarthritis and did not increase cardiovascular outcomes, 46-47

A prophylactically implanted cardioverter defibrillator did not reduce all-cause mortality after a recent myocardial infarction, 58

Renal impairment increased mortality and cardiovascular complications after myocardial infarction, 51

### Myocardial revascularization

An early invasive strategy improved long-term freedom from major adverse cardiac events in older patients with angina, 39

## N

### Naproxen

Lumiracoxib reduced ulcer complications compared with ibuprofen and naproxen in osteoarthritis and did not increase cardiovascular outcomes, 46-47

### Narcotics

Review: Nonsteroidal antiinflammatory drugs provide better pain relief than do opioids for acute renal colic, 42

### Natriuretic peptide, brain

Rapid measurement of B-type natriuretic peptides reduced time to discharge and treatment costs in patients with acute dyspnea (Letter), A-14-A-15 (No. 1)

### Neoplasms

Review: Oral and intravenous antibiotics do not differ for effectiveness in febrile neutropenic patients with cancer, 32

### Neutropenia

Review: Oral and intravenous antibiotics do not differ for effectiveness in febrile neutropenic patients with cancer, 32

### Nicotine

Review: Nicotine replacement therapy is effective in both men and women, 13

## O

### Ofloxacin

A 10-day regimen of levofloxacin was not needed in patients with uncomplicated cellulitis who had an acceptable 5-day response, 45

### Omeprazole

Celecoxib was noninferior to diclofenac plus omeprazole for preventing gastroduodenal ulcer recurrence in high-risk patients with arthritis, 76

Esomeprazole was not better than omeprazole for resolving heartburn in endoscopy-negative reflux disease, 6

### Osteoarthritis

Lumiracoxib reduced ulcer complications compared with ibuprofen and naproxen in osteoarthritis and did not increase cardiovascular outcomes, 46-47

### Osteoporosis

Etidronate, calcium, or both did not reduce fracture rates in patients with asthma receiving glucocorticoid treatment, 44

## P

### Pain

Review: Anticonvulsants are better than placebo for reducing the frequency of migraine attacks, 21

### Patient education

A dietitian-led intervention reduced weight and waist circumference in obese patients with type 2 diabetes, 17

### Peptic ulcer

Celecoxib was noninferior to diclofenac plus omeprazole for preventing gastroduodenal ulcer recurrence in high-risk patients with arthritis, 76

Lumiracoxib reduced ulcer complications compared with ibuprofen and naproxen in osteoarthritis and did not increase cardiovascular outcomes, 46-47

### Peptides

Caspofungin was noninferior to amphotericin B for invasive fungal infections in persistent fever and neutropenia and was better tolerated, 33

Phthalic acids

Picotamide reduced all-cause mortality more than aspirin in type 2 diabetes mellitus and peripheral arterial disease, 61

Polyuria

Review: Estrogen improves symptoms of overactive bladder in postmenopausal women, 48

Postphlebotic syndrome

Below-knee elastic compression stockings reduced development of the postthrombotic syndrome in proximal deep venous thrombosis, 7

Primary health care

*Decision Making in Health and Medicine. Integrating Evidence and Values* (Resource Corner), A-11 (No. 2)

*Screening for Diseases: Prevention in Primary Care* (Resource Corner), A-11 (No. 1)

Primary prevention

*Screening for Diseases: Prevention in Primary Care* (Resource Corner), A-11 (No. 1)

Prostatic neoplasms

Mortality rate from early prostate cancer increased 3-fold after 15 years following the diagnosis, 24

Proton pumps

Review: Misoprostol or COX-2-specific or selective NSAIDs reduce gastrointestinal complications and symptomatic ulcers, 75

Review: Proton-pump inhibitor therapy reduces symptoms in nonulcer dyspepsia better than placebo, 74

Public health

*Evidence-based Public Health* (Resource Corner), A-10 (No. 1)

Receptors, adrenergic,  $\beta$ -2

Gly/Gly and Arg/Arg genotype responses to albuterol differed in mild asthma, 43

R

Renal artery obstruction

Computed tomographic angiography and magnetic resonance angiography were specific, but could not rule out renal artery stenosis, 80

Ribavirin

Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections, 10

Peginterferon  $\alpha$ -2a improved the hepatitis C virologic response in concurrent HIV and chronic hepatitis C virus infections (Letter), A-12-A-13 (No. 1)

S

Sepsis

Review: Glucocorticoids reduced mortality in sepsis in recent (post-1997) but not previous (pre-1989) trials, or all trials combined, 31

Review: Low-dose but not high-dose corticosteroids reduced all-cause mortality in severe sepsis and septic shock, 30

Serotonin reuptake inhibitors

Citalopram did not differ from placebo for treatment of depression in patients > 75 years of age, 79

Shock, septic

Review: Glucocorticoids reduced mortality in sepsis in recent (post-1997) but not previous (pre-1989) trials, or all trials combined, 31

Review: Low-dose but not high-dose corticosteroids reduced all-cause mortality in severe sepsis and septic shock, 30

Simvastatin

Ezetimibe plus simvastatin lowered lipid levels more than simvastatin monotherapy in primary hypercholesterolemia, 16

Smoking cessation

Review: Bupropion and nortriptyline each increase smoking cessation rates, 67

Review: Clonidine is more effective than placebo for long-term smoking cessation but has side effects, 12

Review: Nicotine replacement therapy is effective in both men and women, 13

Stents

Invasive strategy within 24 hours of thrombolysis reduced death, nonfatal reinfarction, and ischemia-induced revascularization in STEMI, 38

Stenting was noninferior to endarterectomy in patients with severe carotid artery stenosis and coexisting conditions, 41

Sulfonamides

Celecoxib was noninferior to diclofenac plus omeprazole for preventing gastroduodenal ulcer recurrence in high-risk patients with arthritis, 76

T

Thromboembolism

Review: Low-molecular-weight heparin reduces recurrent venous thromboembolism better than unfractionated heparin, 71

Thrombolytic therapy

Invasive strategy within 24 hours of thrombolysis reduced death, nonfatal reinfarction, and ischemia-induced revascularization in STEMI, 38

Tomography, x-ray computed

Computed tomographic angiography and magnetic resonance angiography were specific, but could not rule out renal artery stenosis, 80

Trichomonas vaginitis

Review: Vaginal signs and symptoms perform poorly in diagnosing vaginal candidiasis (Correction), A-13 (No. 2)

Review: Vaginal signs and symptoms perform poorly in diagnosing vaginal candidiasis (Letter), A-13 (No. 2)

V

Vascular diseases

Picotamide reduced all-cause mortality more than aspirin in type 2 diabetes mellitus and peripheral arterial disease, 61

Venous thrombosis

Below-knee elastic compression stockings reduced development of the postthrombotic syndrome in proximal deep venous thrombosis, 7

Dalteparin reduced venous thromboembolic events without increased bleeding in acutely ill medical patients, 40

Ventricular dysfunction, left

Review:  $\beta$ -blockers reduced mortality and therapy withdrawal in heart failure but increased dizziness and bradycardia, 36

Ventricular fibrillation

Review: Cardiac resynchronization therapy reduces mortality and hospitalization for heart failure, 35

Verapamil

Trandolapril delayed persistent microalbuminuria in hypertension, type 2 diabetes, and normoalbuminuria, 65

Volunteer workers

Volunteers trained in CPR and the use of automated external defibrillators increased survival after out-of-hospital cardiac arrest, 2

W

Weight loss

Review: Fluoxetine, orlistat, and sibutramine modestly reduce weight in type 2 diabetes, 18

Review: Orlistat and sibutramine are modestly effective for weight loss at 1 year, 19



